

Broad Spectrum Pest Control

TORAC[®] insecticide provides *fast*, broad spectrum control of aphids, Colorado potato beetle, leafhoppers, potato psyllid and thrips in potatoes.

Key Benefits

- Effective contact control on all life stages of pests (eggs, larvae, nymphs, adults)
- Fast mode of action
 - foliar contact insecticide that inhibits cellular respiration in the insect
 - rapid feeding inhibition
 - controls pests usually within 48 hours
- Provides excellent control of 1st and 2nd generation Colorado potato beetle
- Excellent resistance management tool option
 - IRAC Group 21A – Mitochondrial Complex 1 Electron Transport Inhibitors (METI)
 - Ideal rotational chemistry with most commonly used insect control programs

Use Recommendations

- Use Rate: 14 to 21 fl oz/acre, depending on pest
- Application Method: ground, air, chemigation
- Minimum Spray Volume: 20 gallons of water per acre by ground; 5 gallons of water per acre by air. Refer to product label for chemigation recommendations.

Use Information

- Torac is a contact insecticide. Proper application and coverage is critical for maximizing pest control.
- Apply with an adjuvant/spreader to optimize uniformity of coverage and performance.
- Maximum Applications: 2 applications per crop cycle
- Spray Interval: 14 days between applications
- Maximum Dosage: 42.0 fl oz per acre per crop cycle
- Reentry Interval (REI): 12 hours
- Preharvest Interval (PHI): 14 days

Colorado Potato Beetle
Larva on Potato



Untreated



1 day after treatment
Torac @ 21 fl oz/acre