

Winter Weed Control with Venue

| | Chickweed | Sowthistle | Fiddleneck | Groundsel | Willowherb | Prickly Lettuce | Henbit | Stinging Nettle | Mustards | Marestail | Fleabane | Filaree | Malva | Redmaids | Swinecress |
|---|-----------|------------|------------|-----------|------------|-----------------|--------|-----------------|----------|-----------|----------|---------|-------|----------|------------|
| Glyphosate (5lb ai) 2-3 pts | | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Glyphosate (5lb ai) 2-3 pts | | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Glyphosate (5lb ai) 3 pts + Oxyfluorfen 1-2 pts | | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Paraquat 3-4 pts | | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Glufosinate 48-82 fl oz | | | | | | | | | | | | | | | |

Spring and Summer Weed Control with Venue

| | Lambsquarters | Purslane | Field Bindweed | Morningglory | Nightshades | Knotweed | Russian Thistle | Pigweeds | Marestail | Fleabane | Primrose | Wild Radish | Shepherd's-purse | Malva |
|---|---------------|----------|----------------|--------------|-------------|----------|-----------------|----------|-----------|----------|----------|-------------|------------------|-------|
| Glyphosate (5lb ai) 2-3 pts | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Glyphosate (5lb ai) 2-3 pts | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Glyphosate (5lb ai) 2-3 pts + Oxyfluorfen 1-2 pts | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Paraquat 3-4 pts | | | | | | | | | | | | | | |
| Venue 3-4 fl oz + Glufosinate 48-82 fl oz | | | | | | | | | | | | | | |

- For best control, weeds must be at rosette stage or smaller than 4 inches in height.
- For control of large weeds, multiple applications may be needed.
- Increase Glufosinate rates as weeds become larger.

% Control Color Legend

- 90-100% Control
- 80-90% Partial Control
- 60-80% Suppression

Key Benefits

- Broadens the spectrum of weed control
- Increases control of hard-to-kill weeds
- Can be used in newly planted orchards and vineyards
- Can be used for sucker control in a variety of labeled crops
- Reduced risk of volatilization
- 0 day PHI for grapes, olive trees, pome fruits, pomegranates, stone fruits, and tree nuts
- “Caution” signal word makes Venue® easy to use and worker friendly

Mixing Instructions

- Fill spray tank with $\frac{3}{4}$ of the amount of water needed for the intended application; then turn on agitation.
- Follow proper mixing order sequence: water conditioners, wettable powders (WP), water dispersible granules (WDG), liquid flowables (LF), emulsifiable concentrates (EC), and adjuvants and/or oils.
- When tank mixing with glyphosate or paraquat herbicides, always add Venue to the tank *prior* to adding any glyphosate or paraquat products. **Do not add Venue to a tank already containing these products.**
- Always refer to product labels for complete mixing order instructions.

NICHINO
AMERICA®

©2019 Nichino America, Inc. All rights reserved. Venue and Nichino America logo are registered trademarks of Nichino America, Inc. Roundup is a registered trademark of Monsanto Company. Always read and follow all label directions. **888-740-7700** www.nichino.net VEN-TNV-2001

**VENUE**
HERBICIDE

Weed Control on Morningglory



Before Treatment



14 Days After Treatment
Venue @ 3 fl oz + Roundup® @ 32 fl oz